NATIONAL BUREAU OF STANDARDS REPORT

1665

A PROPERTY OF STRONGLY CONTINUOUS PROCESSES

Ъу

Bugene Lukacs



U. S. DEPARTMENT OF COMMERCE NATIONAL BUREAU OF STANDARDS

U. S. DEPARTMENT OF COMMERCE

Charles Sawyer, Secretary

NATIONAL BUREAU OF STANDARDS

A. V. Astin. Acting Director



THE NATIONAL BUREAU OF STANDARDS

The scope of activities of the National Bureau of Standards is suggested in the following listing of the divisions and sections engaged in technical work. In general, each section is engaged in specialized research, development, and engineering in the field indicated by its title. A brief description of the activities, and of the resultant reports and publications, appears on the inside of the back cover of this report.

- 1. Electricity. Resistance Measurements. Inductance and Capacitance. Electrical Instruments. Magnetic Measurements. Electrochemistry.
- 2. Optics and Metrology. Photometry and Colorimetry. Optical Instruments. Photographic Technology. Length. Gage.
- 3. Heat and Power. Temperature Measurements. Thermodynamics. Cryogenics. Engines and Lubrication. Engine Fucls.
- 4. Atomic and Radiation Physics. Spectroscopy. Radiometry. Mass Spectrometry. Physical Electronics. Electron Physics. Atomic Physics. Neutron Measurements. Nuclear Physics. Radioactivity. X-Rays. Betatron. Nucleonic Instrumentation. Radiological Equipment. Atomic Energy Commission Instruments Branch.
- 5. Chemistry. Organic Coatings. Surface Chemistry. Organic Chemistry. Analytical Chemistry. Inorganic Chemistry. Electrodeposition. Gas Chemistry. Physical Chemistry. Thermochemistry. Spectrochemistry. Pure Substances.
- 6. Mechanics. Sound. Mcchanical Instruments. Aerodynamics. Engineering Mechanics. Hydraulics. Mass. Capacity, Density, and Fluid Meters.
- 7. Organic and Fibrous Materials. Rubber. Textiles. Paper. Leather. Testing and Specifications. Organic Plastics. Dental Research.
- 8. Metallurgy. Thermal Metallurgy. Chemical Metallurgy. Mechanical Metallurgy. Corrosion.
- Mineral Products. Porcelain and Pottery. Glass. Refractories. Enameled Metals. Building Stone. Concreting Materials. Constitution and Microstructure. Chemistry of Mineral Products.
- 10. Building Technology. Structural Engineering. Fire Protection. Heating and Air Conditioning. Exterior and Interior Coverings. Codes and Specifications.
- 11. Applied Mathematics. Numerical Analysis. Computation. Statistical Engineering. Machine Development.
- 12. Electronics. Engineering Electronics. Electron Tubes. Electronic Computers. Electronic Instrumentation.
- 13. Ordnance Development. Mechanical Research and Development. Electromechanical Fuzes. Technical Services. Missile Fuzing Research. Missile Fuzing Development. Projectile Fuzes. Ordnance Components. Ordnance Tests. Ordnance Research.
- 14. Radio Propagation. Upper Atmosphere Research. Ionospheric Research. Regular Propagation Services. Frequency Utilization Research. Tropospheric Propagation Research. High Frequency Standards. Microwave Standards.
- 15. Missile Development. Missile Engineering. Missile Dynamics. Missile Intelligence. Missile Instrumentation. Technical Services. Combustion.

May 14, 1952

NBS REPORT

1103-21-5119

1665

A PROPERTY OF STRONGLY CONTINUOUS PROCESSES

рy

Eugene Lukacs



*The preparation of this paper was sponsored by the U. S. Naval

Ordnance Test Station, Inyokern

This report is issuin any form, either: from the Office of the

Approved for public release by the Director of the National Institute of Standards and Technology (NIST) on October 9, 2015

ation, reprinting, or reproduction permission in writing is obtained , Washington 25, D. C.



A PROPERTY OF STRONGLY CONTINUOUS PROCESSES

· by

Augene Lukace National Bureau of Stenderie

1. Introduction.

The purpose of this report is to give a condition which assures that the increments of a stochastic process y(t) are normally distributed.


This condition is of interest in connection with the study of the fundamental random process. The fundamental random process - also called the Wiener process - affords a particularly simple model for certain phenomena. One of the preperties of the fundamental random process is that its increments are normally and independently distributed (for a definition see [4] or [5]). In the present report it is shown that the normality of the increments follows from a certain continuity property.

2. Definitions and statement of the theorem.

In this section we give the definitions necessary to formulate the theorem.

The <u>increment</u> of a stochastic process y(t) over the time interval $(t, t \div 7)$ is the random variable $y(t \div 7) = y(t)$.

A stochastic process is said to be a process with independent increments over non-overlapping time intervals ere completely independent of each other.



A process y(t) is said to be <u>strongly continuous</u> [5] in the interval [a, b] if to every (>0) and y>0 there exists a S=S (<,y) such that for every finite set S of points contained in $\{a,b\}$

(1) P[\$(d, e, B)] = 1 - 7.

Here $\mathcal{E}(\delta,\epsilon,S)$ is the event that the inequalities $|y(t_1)-y(t_k)| \leq \epsilon$ are simultaneously satisfied for all pairs (t_1, t_k) with $|t_1-t_k| < \delta$ belonging to a finite set S of points contained in [a,b].

The symbol P[...] stends here and in the following for the probability of the overt[...] in the brackets.

We are now in a position to formulate the condition manticaed in the introductory section.

Theorem: Lot y(t) be a stochastic process and assume that

(1) y(t) is a process with independent increments


(ii) y(t) is strongly continuous in the interval [2,5].
Then y(b)- y(a) is sormally distributed.

A theorem of this type is due to P. Levy [3] (theorem 16, 3). However it might be of interest to give a different proof using the notation and terminology of H. B. Mann's monograph [5].

3. Ibinichine's theorem.

For the proof we need a theorem of A. Khintchine which gives conditions for the convergence to the normal distribution. This theorem is given in a book by Khintchine, published in the Bussian language. The following formulation was taken from a paper by B. V. Snedenko [1] which is small able in an English translation.

We consider a sequence of sequences X_{nR_1} , X_{nR_2} , ... $X_{nK_{nR_1}}$, n=1,2... at inf) of random variables which are independent within cash esquence. The random variables X_{nR_1} are said to be infinitesimal if for $x_{nR_2} > 0$ the relation $\lim_{n\to\infty} X_n > 0$ the relation $\lim_{n\to\infty} X_n > 0$ uniformly in $X_n > 0$.



We denote by $\Gamma_{n,l}(x)$ the distribution function of the random variable $X_{n,l}(x)$ and state <u>Khimtchine's theorem</u>: If the distributions of the sums (2) $T_n = X_{n,l} + X_{n,l} +$

(3)
$$\lim_{n\to\infty} \sum_{k=1}^{k_n} \int_{|x| \ge 6} dF_{nk}(x) = 0$$
.

For the proof the reader is referred to Gwedenko's paper [1].

4. Proof of the theorem stated in section 2

We consider a sequence $\{S_n\}$ of subdivisions of the interval [a,b].

For the case of simplicity we let S_n be the subdivision $\{f_n\}$. $S_n = (t_0^n, t_1^n, t_2^n, \dots, t_n^n)$ of [a,b] into n equal parts, that is we put $f_j = a + (b-a) j/n$ ($j = 0,1,2,\ldots,n$). We write $f_j = a + (b-a) j/n$ ($j = 0,1,2,\ldots,n$) and show first that the $f_j = f_j = f$

By assumption (ii) the process y(t) is strongly continuous in [a,b]. It is therefore possible to determine for every f > 0 and $\eta > 0$ a $\delta = \delta(\epsilon, \eta)$ such that for every subdivision S_n (4) $P[\delta(\delta, \epsilon, S_n)] \ge 1 - \eta$ We denote by $P[f = X_{n,j} | \delta \epsilon \in j = 1, 2, ..., n]$ the probability that the n inequalities $|X_{n,j}| \le \epsilon \in j = 1, 2, ..., n$ hold simultaneously (*)

probability that all n events occur simultaneously. Thus P[R₁; i= 1,2,...n] means the probability of the simultaneously. Thus P[R₁; i= 1,2,...n] length means that the probability of the simultaneous occurrence of all nevents is at least equal to 1 - n (1=1,2,...n) which means that the probability of the occurrence of each single event is at least equal to 1/2, this statement does not imply anything about the probability of the joint occurrence of the n events.



We next choose a number $f = \mathbb{N}(\xi, n)$ such that $\mathbb{N} \geq \frac{(b-a)}{d(\varepsilon, n)}$.

For any $n \geq N$ the event $C(d, \xi, S_n)$ implies that the N inequalities $\|\mathbf{x}_{n,1}\| \leq \xi(j=1,2,\ldots,n)$ are simultaneously satisfied. We conclude then from (-1) that

- (5) $1 \eta = P[|x_{n_0}| \le \xi; j = 1, 2, ... n]$ and also
- (6) $1 \eta = P[|x_{n_0}|] = \ell]$ for j = 1, 2, ..., if only <math>n = N.

For every $\ell > 0$ and $\eta > 0$ it is therefore possible to find an N = N (ℓ, η) such that for $n \ge N$

- (7) $P[|x_{n,j}| > \xi] \leq \eta$ for j = 1,2,...n.

 This shows that $P[|x_{n,j}| > \xi]$ converges (uniformly in j) to zero as $n \to \infty$, or in other words the $x_{n,j}$ are infinitesimal random variables.

 We consider the sequence of random variables
- (8) $T_n = x_{n,1} + x_{n,2} + \cdots + x_{n,n}$ We have shown that T_n is the sum of infinitesimal random variables which are by assumption independent (within each sequence). Since

 $T_n = \sum_{j=1}^n x_{n,j} = \sum_{j=1}^n [y(t_j^n) - y(t_j^n)] = y(t_n^n) - y(t_n^n) - y(t_n^n) = y(b) - y(a)$ We see that the limiting distribution of the T_n is the distribution of the random variable y(b) - y(a).

We have already shown that it is possible to find for every E>0 and n>0 and n>0

Since the increments $x_{n,j}$ are independently distributed [by assumption (1)] it follows that also

(9) $P[|x_{n,j}| \le \xi; j = 1,2,...k] \ge 1-\eta$ where ξ and η are arbitrary positive numbers and $n \ge 1$ while k is an integer less than n.



For any subdivision Sn of [ab] into h equal parts we introduce the random variable

(10) $M_{abS_n} = M_{ax} [|x_n, 1|, |x_n, 2|, ..., |x_n, n|].$ The statement that the n inequalities $|x_n, j| \le \xi$ for j = 1, 2, ..., nhold simultaneously is equivalent to the statement $M_{abS_n} \le \xi$.

Therefore

P[Mabsn & &] = P[|xn, j| & &; j=1,2,...n] & l-n

and also

(11) P[Mabsn > E] 着 項 ,

for all $\xi > 0$ and $\eta > 0$, provided that $n \ge N$.

We introduce the following n events:

 $B_1^{(n)}$ is the event that the inequality $|x_{n,1}| > \mathcal{E}$ holds.

 $B_j^{(n)}$ (for j=2,3,...n) is the event that the j inequalities

 $|x_{n,1}| \le \mathcal{E}, |x_{n,2}| \le \mathcal{E}, \cdots, |x_{n,j-1}| \le \mathcal{E}, |x_{n,j}| > \mathcal{E} \text{ hold}$ simultaneously. The events $B_1^{(n)}, B_2^{(n)}, \dots, B_n^{(n)}$

are mutually exclusive and exhaust all the cases for which $H_{ab}/m \gg E$. $/(c_{...}S)$ Therefore we see from (11)

(12) $\eta = P[M_{abS_n} > \mathcal{E}] = \sum_{j=1}^{n} P[B_j^{(n)}]$. for all $\mathcal{E} > 0$ and $\eta > 0$, provided that $n \geq N$.

Since the random variables $x_{n,1},\dots,x_{n,n}$ are completely independent we have for $y=2,\dots$

 $P[B_{j}^{n}] = P[1x_{n,k} | \mathcal{E}_{i}; k=1,2,...(j-1)] P[|x_{n,j}| \geq \mathcal{E}_{j}.$

It follows therefore from (9) that

(13) $P[B_j^n] \ge (1-\eta) P[|x_{n,j}| \ge \xi]$ (j=2,...,n)

for any \$ > 0 and n > Oif only n N so that

(14) $\sum_{j=1}^{n} P[B^{n}] \stackrel{\ge}{=} (1-\eta) \stackrel{n}{\sum_{j=1}^{n}} P[|x_{n,j}| > \varepsilon]$ for any $\varepsilon > 0$ and $\eta > 0$ provided that $n \stackrel{\ge}{=} N$.

From (12) and (14) we see that

The state of the s

the second second second second

I Sele, was a special for the

(15)
$$\eta/(1-\eta) = \sum_{j=1}^{n} P[|x_{n,j}| > \xi]$$

for any $\xi > 0, \eta > 0$ if $n \ge N$. Hence we conclude that

$$\lim_{n\to\infty} \sum_{j=1}^{n} P[|x_{n,j}| > \xi] = 0 \quad \text{or}$$

(16)
$$\lim_{n \to \infty} \sum_{j=1}^{n} \int_{|x| > \xi} dF_{n,j}(x) = 0$$

But (16) is exactly Khinchine's condition (3) so that we have shown that the limiting distribution of the T_n , that is the distribution of y (b) - y(a) is normal.




- 1 . B. V. Gmedenko: Limit Theorems for Sums of Independent Random
 Variables. American Mathematical Society Translation No. 45.

 New York (1951). Translated from Uspehi Matem. Nauk 10, 115-165
 (1944).
- 2 . A. Khintchine, Limiting distributions for sums of independent random variables. (in Russian) Moscow Leningrad (1938).
- 3 . Paul Levy Processes stochastiques et mouvement Brownien.

 Gauthiers Villars, Paris 1948
- .4. H. B. Mann Estimation of parameters in certain stochastic processes.

 Sankhya vol. 11, 97-106, (1951)
- 5. H. B. Mann Introduction to the theory of stochastic processes
 depending on a continuous parameter.

 N.B.S. report #1293 (1951) to be published as AMS 24.



THE NATIONAL BUREAU OF STANDARDS

Functions and Activities

The functions of the National Bureau of Standards are set forth in the Act of Congress, March 3, 1901, as amended by Congress in Public Law 619, 1950. These include the development and maintenance of the national standards of measurement and the provision of means and methods for making measurements consistent with these standards; the determination of physical constants and properties of materials; the development of methods and instruments for testing materials, devices, and structures; advisory services to Government Agencies on scientific and technical problems; invention and development of devices to serve special needs of the Government; and the development of standard practices, codes, and specifications. The work includes basic and applied research, development, engineering, instrumentation, testing, evaluation, calibration services and various consultation and information services. A major portion of the Bureau's work is performed for other Government Agencies, particularly the Department of Defense and the Atomic Energy Commission. The scope of activities is suggested by the listing of divisions and sections on the inside of the front cover.

Reports and Publications

The results of the Bureau's work take the form of either actual equipment and devices or published papers and reports. Reports are issued to the sponsoring agency of a particular project or program. Published papers appear either in the Bureau's own series of publications or in the journals of professional and scientific societies. The Bureau itself publishes three monthly periodicals, available from the Government Printing Office: The Journal of Research, which presents complete papers reporting technical investigations; the Technical News Bulletin, which presents summary and preliminary reports on work in progress; and Basic Radio Propagation Predictions, which provides data for determining the best frequencies to use for radio communications throughout the world. There are also five series of nonperiodical publications: The Applied Mathematics Series, Circulars, Handbooks, Building Materials and Structures Reports, and Miscellaneous Publications.

Information on the Bureau's publications can be found in NBS Circular 460, Publications of the National Bureau of Standards (\$1.00). Information on calibration services and fees can be found in NBS Circular 483, Testing by the National Bureau of Standards (25 cents). Both are available from the Government Printing Office. Inquiries regarding the Bureau's reports and publications should be addressed to the Office of Scientific Publications, National Bureau of Standards, Washington 25, D. C.



*